

Cosmic string lens effects on CMB polarization patterns

K. Benabed and F. Bernardeau

Service de Physique Théorique, C.E. de Saclay, 91191 Gif-Sur-Yvette, France.

(February 1, 2008)

Extending the Kaiser-Stebbins mechanism we propose here a method for detecting relics of topological defects such as cosmic strings based on lens-induced small-scale B -type polarization in the cosmic microwave background (CMB). Models of inflation, in which large scale structures of the Universe emerge from the inflaton fluctuations, do not exclude the formation of topological defects at the end of the inflationary phase. In such case, we show that the lens effect of a string on the small scale E -type polarization of the cosmic microwave background induces a significant amount of B -type polarization along the line-of-sight. The amplitude of the effect is estimated for different resolutions of cosmic microwave background experiments.

98.70.Vc, 98.80.Cq, 98.62.Sb

INTRODUCTION

The temperature anisotropies of the cosmic microwave background offer a unique window towards the physics of the early Universe and for the understanding of the large-scale structures. Current observations of the temperature anisotropies power spectrum, C_ℓ , point toward the existence of a well localized first acoustic peak [1]. If this result is confirmed by the next generation of CMB experiments, it supports models of large-scale structure formation from adiabatic scalar fluctuations at the expense of models of topological defects and more particularly of cosmic strings [2] [3]. Furthermore, the shape and normalization of the local matter density power spectrum, $P(k)$, is also in bad agreement with the CMB data for such models [4]. This suggests that *only* a small fraction of the large-scale inhomogeneities might be due to topological defects. However, recent studies have shown that in realistic models of inflation cosmic string formation seems quite natural at the end of the inflationary period: it is a natural outcome in Super-Symmetry inspired scenario [5]; it can also be obtained during a pre-reheating process [6].

The effects of cosmic strings on the last scattering surface temperature map have been described by Kaiser & Stebbins [7]. If a cosmic string is moving against an homogeneous surface of uniform temperature, the energy of the deflected photon [8] is enhanced or reduced (the photons are then blueshifted or redshifted) depending on whether the photon is passing behind or ahead the moving string, a mechanism through which temperature anisotropies are generated. The aim of this paper is to explore the possibilities of having similar effects for the polarization properties. Obviously, if the background surface is unpolarized, the deflected photons remain unpolarized and no effects can be observed. However if the background sky is polarized then the polarization pattern is affected through lens effects and, in particular a geometrical deformation can naturally induce B -type polarization out of the E component. This mechanism has been described for the large-scale structures [9] and

recognized as a major source of B -type small-scale CMB polarization. We are interested here in the case of cosmic strings for which this effect can be easily investigated and visualized. Note that, by doing so, we neglect a possible coupling with an axionic field associated with the string that could induce significant photon-string non-gravitational couplings to a finite distance [10]. The background model of large-scale structure formation of this paper is therefore inflation driven adiabatic fluctuations with a few cosmic strings that may have survived from a late time phase transition although with a significant linear energy density. Note that what we are describing here is a secondary effect from a perturbation theory point of view in the sense that it is quadratic in the metric perturbation: it is a coupling between the local gravitational potential and the potential on the last-scattering surface (the Kaiser-Stebbins effect is also a secondary effect). We do not attempt to describe the primary anisotropies induced at the recombination time that have been examined in other studies [11].

In sect. 1 we examine in detail the effects induced by straight strings and by circular loops. Sect. 2 contains the result of simulations of the effect. Then, in sect. 3, we estimate the detectable amplitude of the B component in the case of a straight string driven deformation.

I. COSMIC STRING LENS EFFECT

In inflationary scenario, at any given scale, scalar perturbations give birth to a scalar polarization pattern – that is to say, to E -type polarization – whereas tensor modes, that can induce both E and B -type polarization, contribute only at very large scale [12]. This result accounts for the symmetry of the fluctuations. It implies that at small scales, the pseudo-scalar B field,

$$B \equiv \Delta^{-1}[(\partial x^2 - \partial y^2)U - 2\partial x \partial y Q], \quad (1)$$

defined* from the Stokes parameters Q and U is zero. The polarization field is then entirely defined by the scalar E field,

$$E \equiv \Delta^{-1}[(\partial x^2 - \partial y^2)Q + 2\partial x\partial y U]. \quad (2)$$

Since the polarization vector is parallel transported along the geodesics, a gravitational lens affects the polarization simply by displacing the apparent position of the polarized light source [14]. In other words, the observed Stokes parameters $\hat{Q}$ and $\hat{U}$ are given in terms of the *primary* (i.e. unlensed) ones by:

$$\hat{Q}(\vec{\alpha}) = Q(\vec{\alpha} + \vec{\xi}), \quad \hat{U}(\vec{\alpha}) = U(\vec{\alpha} + \vec{\xi}). \quad (3)$$

where $\vec{\xi}$ is the displacement field at angular position $\vec{\alpha}$ ($\vec{\alpha}$ is a 2D vector that gives the sky coordinates in the small angle limit). The displacement, $\vec{\xi}$, is given by the integration of the gravitational potential along the line-of-sights. We will assume in this paper that the only potential acting as lens is the cosmic string potential. It obviously depends on the shape, equation of state and dynamics of the string.

Putting (3) in (1), we can write a general expression of $\Delta\hat{B}$ in presence of lenses;

$$\begin{aligned} \Delta\hat{B}(\vec{\alpha}) = & -2Q_{,ij}(\vec{\alpha} + \vec{\xi})(\delta_x^i + \xi_{,x}^i)(\delta_y^j + \xi_{,y}^j) \\ & -2Q_{,i}(\vec{\alpha} + \vec{\xi})\xi_{,xy}^i \\ & +U_{,ij}(\vec{\alpha} + \vec{\xi}) \\ & \times [(\delta_x^i + \xi_{,x}^i)(\delta_y^j + \xi_{,y}^j) - (\delta_y^i + \xi_{,y}^i)(\delta_x^j + \xi_{,x}^j)] \\ & +U_{,i}(\vec{\alpha} + \vec{\xi})(\xi_{,xx}^i - \xi_{,yy}^i). \end{aligned} \quad (4)$$

There are no reasons for the displacement field to preserve a non-zero B -type cosmic microwave background polarization simply because the two scalar field composition (one being the primary scalar perturbation, the other the line-of-sight gravitational potential) breaks the parity invariance.

For illustration we examine here explicitly the two special cases of a straight cosmic string and of a circular cosmic string, both of them in a plane orthogonal to the line-of-sight.

A. The case of a straight string

Let us assume that a straight string is aligned along the y -axis. Then the displacement is uniform at each side of the string. The deflection angle, $\alpha = 4\pi G\mu$ [8] (where G is the Newton constant and μ the string linear energy density) induces a displacement given by,

$$\xi_x = \pm\xi_0, \quad \xi_0 = 4\pi G\mu \frac{\mathcal{D}_{\text{CMB,string}}}{\mathcal{D}_{\text{string}}}, \quad (5)$$

the sign depending on which side of the string one observes; the displacement along y is obviously 0. $\mathcal{D}_{\text{CMB,string}}$ and $\mathcal{D}_{\text{string}}$ are the cosmological angular distances between, respectively, the last scattering surface and the string, and the string and the observer. In the following, we will assume that we are in the most favorable case for detection, when the ratio of the distance is about unity, hence removing any geometrical dependence on the cosmological parameters. Then, the string lays at equal distance between the last scattering surface and the observer[†]. For a $G\mu$ around 10^{-6} [8], the typical expected displacement is about less than 10 arc seconds.

We can write the expression of the Stokes parameters,

$$\begin{aligned} \hat{Q}(x, y) = & Q(x - \xi_0, y)\theta(x - x_0) \\ & + Q(x + \xi_0, y)(1 - \theta(x - x_0)), \end{aligned} \quad (6)$$

where θ is the step function, x_0 is the position of the string. The same expression holds for $\hat{U}$. Since the primary polarization map is B free, the Laplacian of the observable B field is finally given by,

$$\begin{aligned} \Delta\hat{B}(\vec{\alpha}) = & \delta(x - x_0) (|U_{,x}|^+ - 2|Q_{,y}|^+) \\ & + \delta'(x - x_0) |U|^+, \end{aligned} \quad (7)$$

where we define

$$|X|^+ \equiv \hat{X}(x_0^+) - \hat{X}(x_0^-) = X(x_0 - \xi_0) - X(x_0 + \xi_0). \quad (8)$$

One can note in Eq. (7) that the effect is entirely due to the discontinuity induced by the string on the polarization map. Furthermore, the B component of the polarization will only be non-zero on the string itself. Obviously, the efficiency with which such an effect will be observed depends on the angular precision of the detectors as we discuss later.

B. The case of a circular string

The case of a collapsing circular string loop allows to complement our analysis with the effect of the string curvature. As shown in [15] the lens effect of such a string, when facing the observer, is equivalent to the one of a static linear mass distribution. The structure of the displacement field is then simple. Let us consider a loop centered at the origin of our coordinate and of radius α_l . If one observes towards a direction through the loop, the

*Throughout the paper we work in the small angular scale limit. See [13] for a general discussion of these properties.

[†]this means a redshift of 3 for an Einstein-de Sitter universe.

displacement is nil. Outside the loop, the displacement decreases as α_l/α . We have

$$\vec{\xi}(\vec{\alpha}) = -2\xi_0 \frac{\alpha_l}{\alpha^2} \vec{\alpha} \quad \text{for } \alpha > \alpha_l. \quad (9)$$

Note that in ξ_0 , μ is an effective quantity that contains the effects of dynamics as well. Then, $\hat{Q}$ is

$$\begin{aligned} \hat{Q}(\vec{\alpha}) = & Q \left[\vec{\alpha} \left(1 - 2\xi_0 \frac{\alpha_l}{\alpha^2} \right) \right] \theta(\alpha - \alpha_l) \\ & + Q(\vec{\alpha}) [1 - \theta(\alpha - \alpha_l)]. \end{aligned} \quad (10)$$

Two effects are induced in this case. The first one, also present in the straight string case, comes from the discontinuity of the polarization field; this is the *strong lensing* effect. It is due to the existence of a critical region in the source plane where objects can have multiple images (two in this case, but it can be more in general [15]). The second one is a *weak lensing* effect simply due the deformation of the source plane; it will be small compared to the other. This latter effect is investigated in more detail in [16]. We expect these two effects to be present for any string model.

II. SIMULATED MAPS

We present in Figs. 1-4 the simulation results of circular cosmic string (30 arc minutes radius, $\xi_0 = 5''$). The cosmic microwave background polarization realization uses C_ℓ calculated with a standard Λ CDM model. Only scalar primary perturbations are used here since we do not expect any significant tensor mode at such a small scale; without the string, there is no signal in the B component.

The hot and cold (black and white) patches run along the string path. They come from the δ' term in the eq. (7). Its amplitude is the result of a finite difference in the U field at distance $2\xi_0$. We will see in last section that, at small filtering resolution, this term dominates the amplitude of B polarization.

It is interesting to note here that the *weak lensing* effect is negligible at these scales. Besides, even the discontinuity effect is really small at a 5 arc minute angular scale. A 4 times better resolution enhances significantly the signal (about 10 times). Note also that the hot and cold spots along the strings have the same linear sizes as the typical peaks of the polarization field. We expect that this feature and the very peculiar shape of the effect on the cosmic microwave background polarization maps could help discriminating between this effect and other secondary polarization sources or foregrounds (lensing from large-scale structures, dust polarization...) The extraction of a clean cosmic microwave background polarization out of a signal with foregrounds has been studied [19] at larger scale. Few is known about contamination of the B signal at the scale we are looking at here.



FIG. 1. B field for a circular loop crossing a $50' \times 50'$ window. The filter resolution is 5 arc minutes. At this scale, the B field is less than 1% the E one. The very faint patches that can be noticed above the string are *weak lensing* effect signature (a few percent of the *strong lensing* effect coming from the critical region).



FIG. 2. Same as Fig. 1 for the E field. At this scale the string remains completely "diluted" in the E field and cannot be seen (the effect is smaller than 1% of the mean primary E signal).

III. AMPLITUDE OF A STRAIGHT STRING EFFECT

We come back to the case of a straight cosmic string. It is easy to estimate the amplitude of the effect which only consists of a discontinuity. At small angles we can decompose the E field in plane wave Fourier modes,

$$E(\vec{\alpha}) = \int \frac{d^2l}{2\pi} \tilde{E}(\vec{l}) e^{i\vec{\alpha} \cdot \vec{l}}. \quad (11)$$

It is straightforward to write for B , in Fourier space,

$$\begin{aligned} \Delta \hat{B}(x, y) &= 2 \int \frac{d^2l}{2\pi} \tilde{E}(\vec{l}) e^{i\vec{\alpha} \cdot \vec{l}} (e^{i\xi_0 l_x} - e^{-i\xi_0 l_x}) \\ &\times \left[\frac{l_x l_y}{l^2} \delta'(x - x_0) + i \frac{l_y^3}{l^2} \delta(x - x_0) \right]. \end{aligned} \quad (12)$$

This expression makes sense only if convolved with a *test function*; that is to say convolved with a suitable window function. For simplicity, we assume that our observational device window is described by a Gaussian window function W of width α_w ,

$$W(\vec{\alpha}) = \frac{1}{2\pi\alpha_w} e^{-\frac{\alpha^2}{2\alpha_w^2}}, \quad \tilde{W}(k_x, k_y) = e^{-\frac{\alpha_w^2 (k_x^2 + k_y^2)}{2}}. \quad (13)$$

Then we have,

$$\begin{aligned} \Delta \hat{B}_W(x, y) &= 2 \int \frac{d^2l}{2\pi} \frac{dk}{2\pi} \tilde{E}(\vec{l}) e^{i[x_0(l_x - k) + xk + y l_y]} \\ &\times \tilde{W}(k, l_y) (e^{i\xi_0 l_x} - e^{-i\xi_0 l_x}) \left[i \frac{l_x l_y k}{l^2} + i \frac{l_y^3}{l^2} \right]. \end{aligned} \quad (14)$$

The r.m.s. of $\Delta \hat{B}_W$ can then be expressed as a function of the E power spectrum $C_E(l)$,

$$\begin{aligned} \left\langle (\Delta \hat{B}_W)^2 \right\rangle &= 2e^{-\frac{(x-x_0)^2}{\alpha_w^2}} \int \frac{d^2l}{\pi^3} C_E(l) \sin^2(\xi_0 l_x) \\ &\times e^{-l_y^2 \alpha_w^2} \left[\frac{l_x^2 l_y^2 (x - x_0)^2}{l^4 \alpha_w^6} + \frac{l_y^6}{l^4 \alpha_w^2} \right]. \end{aligned} \quad (15)$$

$C_E(l)$, has a natural cutoff due to the Silk damping scale, l_{damp} ($1/l_{\text{damp}} \sim 10'$), a scale much bigger than the induced displacement. Therefore, we can replace $\sin^2(\xi_0 l_x)$ by its expansion $\xi_0^2 l_x^2$. Then the amplitude of the effect grows like ξ_0 . Besides, if the size of the window is smaller than the typical scale of ΔE structures, $\alpha_w \ll \alpha_{\text{peaks}}$ with $\alpha_{\text{peaks}} \sim 10^{-3}$, we have $\exp(-l_y^2 \alpha_w^2) \sim 1$. And from Eq. (15) we can then calculate:

$$\frac{\langle (\Delta \hat{B}_w(x=x_0 \pm \alpha_w, y))^2 \rangle^{1/2}}{\langle (\Delta E)^2 \rangle^{1/2}} = \frac{1}{4\sqrt{\pi e}} \frac{\xi_0}{\alpha_w} \sqrt{5 + 8 \frac{\alpha_{\text{peaks}}^2}{\alpha_w^2}}. \quad (16)$$

The distance to the string, $x = x_0 \pm \alpha_w$, have been chosen to give a realistic account of the effect (in the simulation



FIG. 3. Same as 1 with a better resolution (1.2'). The discontinuity effect is less diluted. The B field is now in amplitude about 10% of the typical E fluctuations.



FIG. 4. Same as 2 for the 1.2' resolution. The discontinuity effect is now visible in E . Lens effects induce mode couplings that create structures at very small scales dominating over primary structures.




FIG. 5. A comparison of the exact computation of $\sqrt{\langle \Delta B^2 \rangle} / \sqrt{\langle \Delta E^2 \rangle}$ (dashed line) at $x = x_0 \pm \alpha_w$ and $\xi_0 = 10''$ and its approximation (see eq. 16). The amplitude of the effect is about 10% at 1' scale, about 1.5% at 5'. The agreement between the exact amplitude and eq. (16) weakens for scales above $2 \sim 3'$.

this correspond almost to the peak of the hot and cold patches). Fig.5 shows these results for a Λ CDM model – the only dependence with the cosmological parameters appears in the position of the polarization peaks which depends essentially on the global curvature of the Universe. Our approximation is not exact at 5', but is fairly accurate at 1'. Numerically, we found that, at 5' resolution, the amplitude of the effect evolves like $\sim 325 \xi_0$ at the position $x = x_0 \pm \alpha_w$; it is $\sim 1.5 \times 10^{-2}$ when $\xi_0 = 10''$. The slope at small α_w is due to the α_w^2 / α_w^2 in eq. (16) that is to say to the δ' in eq. (7). This is in good agreement with the simulations that suggested an effect of this order of magnitude dominated, at small α_w , by the finite difference in U field at $2\xi_0$ scale.

CONCLUSIONS

The so far the planned cosmic microwave background experiments will have, at their very best, a 5' resolution [17]. We showed that at this angular scale and assuming that it exists a string with $\xi_0 = 10''$ (which is perhaps somewhat enthusiastic), we can expect a signal in B -type polarization with an amplitude of about 1% the signal in E -type polarization. This signal is too weak to be actually detected. Beside, the weak lensing is expected to induce B -type polarization at the same scale and is likely to hide any string effect. Improving the resolution of the detector however will dramatically enhance the detectability of a string effect; at 1' scale, we expect indeed to gain a factor 10 in the amplitude of the effect that should make the detection possible. The detection of cosmic strings through this effect (or through the Kaiser-Stebbins effect which also requires a good angular resolution) will probably be possible only with the

post Planck generation of instruments [18].

ACKNOWLEDGMENTS

The authors would like to thank P. Peter, L. Kofman and especially J.P. Uzan for encouraging discussions and comments on the manuscript. We also thank A. Riazuelo for the use of his Boltzmann CMB code.

-
- [1] C. B. Netterfield, M. J. Devlin, N. Jarosik, L. Page, E. J. Wollack, *ApJ* **474**, 47 (1997); A. de Oliveira-Costa et al., *ApJL* **509** (1998); K. Coble et al. *astro-ph/9902195*
 - [2] T.W. Kibble, *J. Phys.* **A9**, 1387 (1976).
 - [3] A. Albrecht, R. A. Battye, J. Robinson, *Phys. Rev.* **D59**, 023508 (1999); R. Durrer, M. Kunz, A. Melchiorri, *astro-ph/9811174*
 - [4] A. Albrecht, R. A. Battye, J. Robinson, *Phys. Rev. Lett.* **79** 4736-4739 (1997).
 - [5] P. Binetruy, C. Deffayet, E. Dudas, P. Ramond, *Phys.Lett.* **B441** 163-172 (1998); P. Binetruy, C. Deffayet, P. Peter, *Phys. Lett.* **B441** 52-59 (1998); R. Jeanerot, *Phys. Rev.* **D56**, 6205-6216 (1999).
 - [6] I. Tkachev, S. Khlebnikov, L. Kofman & A. Linde, *Phys. Lett.* **B440**, 262-268 (1998).
 - [7] N. Kaiser & A. Stebbins, *Nature*, **310**, 391-393, (1984).
 - [8] A. Vilenkin & P. Shellard, “*Cosmic Strings and other Topological Defects*”, Cambridge monograph on mathematical physics, Cambridge University Press, England, 1994.
 - [9] M. Zaldarriaga & U. Seljak, *Phys. Rev.* **D58**: 023003 (1998)
 - [10] J.A. Harvey & S.G. Naculich, *Phys.Lett.* **B217** 231-237 (1989)
 - [11] U. Seljak, U. Pen & N. Turok, *Phys. Rev. Lett.* **79** 1615-1618 (1997)
 - [12] U. Seljak, *ApJ* **482**, 6, (1997); W. Hu & M. White, *New Astronomy* **2**, 323 (1997); J. Lesgourgues, D. Polarski, S. Prunet, A.A. Starobinsky, *gr-qc/9906098*
 - [13] M. Zaldarriaga & U. Seljak, *Phys. Rev.* **D55**, 1830-1840 (1997); M. Kamionkowski, A. Kosowsky, A. Stebbins, *Phys.Rev.* **D55** 7368-7388 (1997).
 - [14] P. Schneider, J. Ehlers & E.E. Falco, “*Gravitational lenses*”, Springer-Verlag, 1992. V. Faraoni, *Astron. Astrophys.* **272** ,385-388 (1993).
 - [15] A.A. de Laix & T. Vachaspati, *Phys.Rev.* **D54**, 470-4791, (1996).
 - [16] K. Benabed & F. Bernardeau (in preparation)
 - [17] phase A report study for the Planck mission, <http://astro.estec.esa.nl/SA-general/Projects/Planck/report/report.html>
 - [18] J.B. Peterson et al. *astro-ph/9907276*
 - [19] F.R. Bouchet, S. Prunet & S. K. Sethi *MNRAS* **302**,663 (1999) ; S. Prunet, S.K. Sethi & F.R. Bouchet *astro-ph/9911243*